



Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines

Makarov, Ilya; Fristrup, Peter; Madsen, Robert

Publication date:
2010

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Makarov, I., Fristrup, P., & Madsen, R. (2010). *Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines*. Abstract from 8th European Conference on Computational Chemistry , Lund, Sweden.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Mechanistic study of the ruthenium-catalyzed synthesis of amides from alcohols and amines

Ilya Makarov, Peter Fristrup, and Robert Madsen

Technical University of Denmark, DK-2800 Lyngby, Denmark

The amide bond is one of the most important linkages in organic chemistry and constitutes the key functional group in peptides, polymers and many pharmaceuticals.^{1,2} Amides are often prepared by coupling of carboxylic acids and amines by the use of either a coupling reagent³ or by prior conversion of the carboxylic acid into a derivative such as the acid chloride. Recently in Madsen group a new method for amide synthesis has been discovered where alcohols and amines are coupled directly with the liberation of hydrogen⁵ (Scheme 1).



Scheme 1. Amide formation from alcohols and amines.

The mechanistic studies of this reaction are divided into an experimental part and a theoretical part. In the experimental studies reaction orders for all reactants are determined, para-substituted benzyl alcohols are used as substrates to elucidate the electronic influence of the para substituent, *i.e.* a Hammett study. To gain further insight into the mechanism, a computational study is also carried out. By using DFT calculations with the B3LYP or M06 functionals Gibbs free energies and structures are determined for some metal complexes and transition states in the proposed catalytic cycle. From the experimental and theoretical studies the rate determining step will be established which is important to further optimize the reaction. With an accurate theoretical model, it will be possible to design improved catalyst structures *in silico*, e.g. to determine the optimum *N*-heterocyclic carbene ligand for the reaction.

1. a) J. W. Bode, *Curr. Opin. Drug Discov. Dev.* **2006**, 9, 765-775; b) K. E. Gonsalves, P. M. Mungara, *Trends Polym. Sci.* **1996**, 4, 25-31.
2. Of the 53 small molecule drugs in 2003 with an annual sale of more than \$1 billion, 9 contained an amide, see J. S. Carey, D. Laffan, C. Thomson, M. T. Williams, *Org. Biomol. Chem.* **2006**, 4, 2337-2347.
3. S.-Y. Han, Y.-A. Kim, *Tetrahedron* **2004**, 60, 2447-2467.
4. C. A. G. N. Montalbetti, V. Falque, *Tetrahedron* **2005**, 61, 10827-10852.
5. L. U. Nordstrøm, H. Vogt, R. Madsen, *J. Am Chem. Soc.* **2008**, 130, 17672-17673.